arXiv:1407.6515v1 [math.GN] 24 Jul 2014

ON GIBSON FUNCTIONS WITH CONNECTED GRAPHS

OLENA KARLOVA* AND VOLODYMYR MYKHAYLYUK**

Abstract. A function $f : X \to Y$ between topological spaces is said to be a *weakly Gibson function* if $f(\overline{G}) \subseteq \overline{f(G)}$ for any open connected set $G \subseteq X$. We call a function $f : X \to Y$ *segmentary connected* if X is topological vector space and $f([a, b])$ is connected for every segment $[a, b] \subseteq X$. We show that if X is a hereditarily Baire space, Y is a metric space, $f : X \to Y$ is a Baire-one function and one of the following conditions holds: (i) X is a connected and locally connected space and f is a weakly Gibson function, (ii) X is an arcwise connected space and f is a Darboux function, (iii) X is a topological vector space and f is a segmentary connected function, then f has a connected graph.

1. Introduction

In 1907, J. Young showed [16] that for a Baire-one function $f : \mathbb{R} \to \mathbb{R}$, the Darboux property is equivalent to the following property: for every $x \in \mathbb{R}$ there exist $(x_n)_{n=1}^{\infty}$ and $(y_n)_{n=1}^{\infty}$ such that $x_n \uparrow x$, $y_n \downarrow x$ and $\lim\limits_{n\to\infty} f(x_n) = \lim\limits_{n\to\infty} f(y_n) = f(x)$. In more general spaces, functions having the property of Young are said to be peripherally continuous (see definition in Section 2).

K. Kuratowski and W. Sierpiński in 1922 proved that a Baire-one function $f : \mathbb{R} \to \mathbb{R}$ has the Darboux property if and only if f has a connected graph [12]. Remark that for functions $f : \mathbb{R} \to Y$, where Y is a topological space, having a connected graph is equivalent to being a connectivity function (see definition in Section 2).

M. Hagan [10] and G. Whyburn [15] showed that a function $f : \mathbb{R}^n \to \mathbb{R}$ for $n > 1$ is connectivity if and only if f is peripherally continuous.

J. Farková in [8] introduced a similar property to the Young property: a function $f : X \to Y$ between metric spaces X and Y has the property $\mathcal{A}$ if for any open connected non-trivial set $G \subseteq X$ and a point $x \in \overline{G}$ there exists a sequence $(x_n)_{n=1}^{\infty}$, $x_n \in G \setminus \{x\}$, such that $\lim\limits_{n\to\infty} x_n = x$ and $\lim\limits_{n\to\infty} f(x_n) = f(x)$.

2010 *Mathematics Subject Classification.* Primary 26B05, 54C08; Secondary 26A21, 54C10.

Key words and phrases. Gibson function, Darboux function, peripherally continuous functions, function with connected graph, Baire-one function.

She proved that for $X = \mathbb{R}$ every Darboux function has the property $\mathcal{A}$, but for $X = \mathbb{R}^2$ this is not true. Moreover, Farková showed that if X is a connected and locally connected complete metric space, Y is a separable metric space and $f : X \to Y$ is a Baire-one function having the property $\mathcal{A}$, then f has a connected graph.

Recently, K. Kellum [11] introduced and investigated properties of Gibson and weakly Gibson real-valued functions (see Definition 3.1) defined on a connected space. Some properties of Gibson functions (in particular, Darboux-like properties or continuity of Baire-one Gibson functions) are also studied by M.J. Evans and P.D. Humke in [7].

In this work we introduce a strong Gibson property (see Definition 3.1) which is equivalent to the property $\mathcal{A}$ for a function between metric spaces. In Section 3 we study how weak and strong Gibson properties connect with the property of peripherally continuity and with the Darboux property. In the forth section we consider Baire-one Gibson functions. In Section 5 we introduced segmentary connected functions defined on a topological vector space and investigate some of their properties. Finally, in the sixth section we prove different generalizations of the mentioned result from [8]. We show that if X is a hereditarily Baire space, Y is a metric space and $f : X \to Y$ is a Bare-one function, then each of the following conditions is sufficient for f to have a connected graph: (i) X is a connected and locally connected space and f is weakly Gibson; (ii) X is an arcwise connected space and f is a Darboux function; (iii) X is a topological vector space and f is a segmentary connected function.

2. Preliminaries

We will denote by $\mathcal{T}(X)$ and $\mathcal{C}(X)$ the collection of all open and connected subsets of a topological space X, respectively. Let $\mathcal{G}(X) = \mathcal{T}(X) \cap \mathcal{C}(X)$.

Let X and Y be topological spaces and let $f : X \to Y$ be a function. Then

- f is a *Darboux function*, $f \in \mathrm{D}(X, Y)$, if $f(C)$ is connected for any $C \in \mathcal{C}(X)$;
- f is a *weakly Darboux function*, $f \in \mathrm{wD}(X, Y)$, if $f(G)$ is connected for any $G \in \mathcal{G}(X)$;
- f *has a connected graph*, $f \in \mathrm{CG}(X, Y)$, if the set $\mathrm{Gr}\,(f) = \{(x, f(x)) : x \in X\}$ is connected;
- f is a *connectivity function*, $f \in \mathrm{Conn}(X, Y)$, if the graph of $f|_C$ is connected for any $C \in \mathcal{C}(X)$;
- f is *peripherally continuous at* $x \in X$ if for an arbitrary open neighborhoods U and V of x and $f(x)$, respectively, there exists an open neighborhood W of x such that $W \subseteq U$ and $f(\mathrm{fr}\, W) \subseteq V$, where $\mathrm{fr}\, W$ denotes the boundary of W; f is *peripherally continuous on* X, $f \in \mathrm{PC}(X, Y)$, if f is peripherally continuous at every point x of X;

- f is a *Baire-one function*, $f \in \mathrm{B}_1(X,Y)$, if there exists a sequence of continuous functions $f_n : X \to Y$ such that $\lim\limits_{n\to\infty} f_n(x) = f(x)$ for every $x \in X$;
- f is *pointwise discontinuous* if the set of all continuity points of f is dense in X;
- f is *barely continuous* if the restriction $f|_F$ has a continuity point for any closed set $F \subseteq X$.

The class of all continuous functions between X and Y we denote by $\mathrm{C}(X,Y)$.

It is easy to see that

$$\mathrm{C}(X,Y) \subseteq \mathrm{Conn}(X,Y) \subseteq \mathrm{D}(X,Y) \subseteq \mathrm{wD}(X,Y)$$

for arbitrary topological spaces X and Y.

The basic relations between the classes described above for $X = Y = \mathbb{R}$ are given in the following chart.

$$\mathrm{C}(\mathbb{R},\mathbb{R}) \subset \mathrm{Conn}(\mathbb{R},\mathbb{R}) = \mathrm{CG}(\mathbb{R},\mathbb{R}) \subset \mathrm{D}(\mathbb{R},\mathbb{R}) \subset \mathrm{PC}(\mathbb{R},\mathbb{R}).$$

For the Baire-one functions the chart changes to the following.

$$\mathrm{C}(\mathbb{R},\mathbb{R}) \subset \mathrm{Conn}(\mathbb{R},\mathbb{R}) = \mathrm{CG}(\mathbb{R},\mathbb{R}) = \mathrm{D}(\mathbb{R},\mathbb{R}) = \mathrm{PC}(\mathbb{R},\mathbb{R}).$$

If $X = \mathbb{R}^n$, $Y = \mathbb{R}$, $n \geq 2$, then

$$\mathrm{C}(\mathbb{R}^n,\mathbb{R}) \subset \mathrm{Conn}(\mathbb{R}^n,\mathbb{R}) = \mathrm{PC}(\mathbb{R}^n,\mathbb{R}) \subset \mathrm{D}(\mathbb{R}^n,\mathbb{R}).$$

For a thorough treatment we refer the reader to [5, 4, 9].

Topological space X is *hereditarily Baire* if every closed subset of X is a Baire space.

Topological space X is *resolvable* if X is a union of two disjoint everywhere dense subsets.

A subset A of topological vector space X is *balanced* if $\lambda A \subseteq A$ whenever $|\lambda| \leq 1$.

For a function $f : X \to Y$ between topological spaces by $C(f)$ and $D(f)$ we denote the sets of continuity and discontinuity points of f, respectively.

3. The Gibson properties and their connection with the Darboux-like properties

Definition 3.1. Let X and Y be topological spaces. A function $f : X \to Y$ is

- *weakly Gibson at* $x \in X$ if $f(x) \in \overline{f(G)}$ for any $G \in \mathcal{G}(X)$ such that $x \in \overline{G}$;
- *Gibson at* $x \in X$ if $f(x) \in \overline{f(G)}$ for any $G \in \mathcal{T}(X)$ such that $x \in \overline{G}$;
- *strongly Gibson at* $x \in X$ if $f(x) \in \overline{f(G \cap U)}$ for any $G \in \mathcal{G}(X)$ such that $x \in \overline{G}$ and for any open neighborhood U of x.

A function $f : X \to Y$ has *(weak, strong) Gibson property on* X if it has such a property at every point of X. Classes of all such functions we denote by $\mathrm{wG}(X, Y)$, $\mathrm{G}(X, Y)$ and $\mathrm{sG}(X, Y)$, respectively.

It is evident that

$$\mathrm{C}(X, Y) \subseteq \mathrm{sG}(X, Y) \subseteq \mathrm{wG}(X, Y)$$

for any topological spaces X and Y.

Proposition 3.2. *Let X be a locally connected space such that $\mathcal{G}(X)$ is closed under finite intersections and Y be a topological space. Then*

$$\mathrm{wG}(X, Y) = \mathrm{sG}(X, Y).$$

Proof. Let $f \in \mathrm{wG}(X, Y)$, $x \in X$, $G \in \mathcal{G}(X)$, $x \in \overline{G}$ and U be a neighborhood of x. We can choose an open connected neighborhood V of x such that $V \subseteq U$. Then $V \cap G \in \mathcal{G}(X)$ and $x \in \overline{V \cap G}$. Therefore, $f(x) \in \overline{f(V \cap G)} \subseteq \overline{f(U \cap G)}$. Hence, $f \in \mathrm{sG}(X, Y)$. □

Corollary 3.3. $\mathrm{wG}(\mathbb{R}, Y) = \mathrm{sG}(\mathbb{R}, Y)$ *for an arbitrary Y.*

Proposition 3.4. *Let X be a topological space and Y be a T_1-space. Then*

$$\mathrm{D}(X, Y) \subseteq \mathrm{wG}(X, Y).$$

Proof. Let $f \in \mathrm{D}(X, Y)$. Choose $x \in X$ and $G \in \mathcal{G}(X)$ such that $x \in \mathrm{fr}\, G$. Fix an arbitrary neighborhood V of $f(x)$ in Y. Since G is connected, $G \cup \{x\}$ is connected too. Then $f(G \cup \{x\})$ is connected since f has the Darboux property. Suppose that $f(G) \subseteq Y \setminus V$. Then

$$f(G \cup \{x\}) = f(G) \cup f(x) \subseteq (Y \setminus V) \sqcup \{f(x)\},$$

which contradicts the fact that $f(G \cup \{x\})$ is connected. □

We will apply the following results to construct a function $f : \mathbb{R}^2 \to \mathbb{R}$ of the first Baire class such that f has the weak Gibson property, but has no strong Gibson property.

Proposition 3.5. *Let X, Y and Z be topological spaces, $f \in \mathrm{wG}(X, Z)$ and $g(x, y) = f(x)$ for all $(x, y) \in X \times Y$. Then $g \in \mathrm{wG}(X \times Y, Z)$.*

Proof. Fix $G \in \mathcal{G}(X \times Y)$ and $p_0 = (x_0, y_0) \in X \times Y$, $p_0 \in \overline{G}$. Choose any neighborhood W of $g(p_0)$ in Z. Since a projection is an open map, $U = \mathrm{pr}_{\mathbb{R}} G$ is an open and connected set in X. Notice that $x_0 \in \overline{U}$. Since f has the weak Gibson property at x_0, there exists $x \in U$ such that $f(x) \in W$. Take an arbitrary $(x, y) \in (\{x\} \times Y) \cap G$. Then $g(x, y) = f(x) \in W$. Hence, g has is weakly Gibson at p_0. □

Proposition 3.6. *Let $f : \mathbb{R} \to \mathbb{R}$ be a function and $g(x, y) = f(x)$ for all $(x, y) \in \mathbb{R}^2$. If $g \in \mathrm{sG}(\mathbb{R}^2, \mathbb{R}) \cap \mathrm{B}_1(\mathbb{R}^2, \mathbb{R})$, then $f \in \mathrm{C}(\mathbb{R}, \mathbb{R})$.*

Proof. Suppose the contrary and choose $x_0 \in D(f)$. Then there exists such a sequence $(x_n)_{n=1}^{\infty}$ that $\lim\limits_{n\to\infty} x_n = x_0$ and $\lim\limits_{n\to\infty} f(x_n) \neq f(x_0)$. Without loss of the generality we may assume that $x_n > x_0$ for all $n \in \mathbb{N}$. Remark that the sets $\Gamma_1 = \mathrm{Gr}(f|_{[x_0,+\infty]})$ and $\Gamma_2 = \mathrm{Gr}(f|_{(x_0,+\infty]})$ are connected since $f : \mathbb{R} \to \mathbb{R}$ is a Baire-one Darboux function. For every $p = (x, f(x)) \in \Gamma_2$ write $B_p = B(p, \frac{|x-x_0|}{2})$. Let $G = \bigcup\limits_{p\in\Gamma_2} B_p$. It is easy to see that there exists $y_0 \in \mathbb{R}$ such that $p_0 = (x_0, y_0) \in \overline{G}$ and $y_0 \neq f(x_0)$. Then g does not have strong Gibson property at p_0. Indeed, if $(p_n)_{n=1}^{\infty}$ be a sequence of points $p_n = (x'_n, y_n) \in G$ with $p_n \to p_0$, then $g(p_n) = f(x'_n) \to y_0 \neq f(x_0) = g(p_0)$, which implies a contradiction. □

Propositions 3.5 and 3.6 now leads to the following example.

Example 3.7. *Let $f : \mathbb{R} \to \mathbb{R}$, $f(x) = \sin\frac{1}{x}$ if $x \neq 0$ and $f(x) = 1$ if $x = 0$, and let $g : \mathbb{R}^2 \to \mathbb{R}$, $g(x, y) = f(x)$. Then $g \in \mathrm{wG}(\mathbb{R}^2, \mathbb{R}) \setminus \mathrm{sG}(\mathbb{R}^2, \mathbb{R})$.*

The following result shows that a strongly Gibson function $f : X \to Y$ can be everywhere discontinuous even in the case where $X = \mathbb{R}$.

Proposition 3.8. *Let X and Y be topological spaces, $X = A \sqcup B$, where A and B are dense in X, $y_1, y_2 \in Y$, $y_1 \neq y_2$ and $f : X \to Y$ be such a function that $f(x) = y_1$ if $x \in A$, and $f(x) = y_2$ if $x \in B$. Then $f \in \mathrm{sG}(X, Y)$.*

Proof. Fix $x \in X$ and $G \in \mathcal{G}(X)$ such that $x \in \mathrm{fr}\, G$. Choose an arbitrary open neighborhoods U and V of x and $f(x)$ in X and Y, respectively. Since $\overline{A} = \overline{B} = X$ and $U \cap G$ is open set, there exist $x' \in A \cap U \cap G$ and $x'' \in B \cap U \cap G$. Then either $f(x') \in V$, or $f(x'') \in V$. Therefore, $f(x) \in \overline{f(U \cap G)}$. □

Proposition 3.9. *Let X be a connected resolvable space, Y be such a T_1-space that $|Y| \geq 2$. Then*

$$\mathrm{sG}(X, Y) \setminus \mathrm{D}(X, Y) \neq \emptyset.$$

Proof. Let X_1 and X_2 are disjoint dense sets in X such that $X = X_1 \cup X_2$. Choose distinct points $y_1, y_2 \in Y$ and consider a function $f : X \to Y$, $f(x) = y_1$ if $x \in X_1$, and $f(x) = y_2$ if $x \in X_2$. According to Proposition 3.8, $f \in \mathrm{sG}(X, Y)$. But $f(X) = \{y_1, y_2\}$ is not connected, then $f \notin \mathrm{D}(X, Y)$. □

The following lemma will be useful.

Lemma 3.10. [14, p. 136] *Let A and B be such subsets of topological space X that A is connected, $A \cap B \neq \emptyset$ and $A \setminus B \neq \emptyset$. Then $A \cap \mathrm{fr} B \neq \emptyset$.*

Proposition 3.11. *Let X be a regular T_1-space and Y be a topological space. Then*

$$\mathrm{PC}(X, Y) \subseteq \mathrm{sG}(X, Y).$$

Proof. Let $f \in \mathrm{PC}(X,Y)$, $G \in \mathcal{G}(X)$, $x \in \mathrm{fr}G$. Fix any neighborhoods U and V of x and $f(x)$ in X and Y, respectively. Choose an arbitrary $x_0 \in G$. Since X is regular T_1-space, there exists an open neighborhood U_1 of x such that $\overline{U}_1 \subseteq U$ and $x_0 \notin U_1$. By peripherally continuity of f at x, there exists such an open neighborhood W of x that $W \subseteq U_1$ and $f(\mathrm{fr}\, W) \subseteq V$. Obviously, $G \cap W \neq \emptyset$ and $G \setminus W \neq \emptyset$. Then Lemma 3.10 implies that $G \cap \mathrm{fr}W \neq \emptyset$. Take $x' \in G \cap \mathrm{fr}W$. Then $x' \in G \cap U$ and $f(x') \in V$. Therefore, $f \in \mathrm{sG}(X,Y)$. □

Lemma 3.12. *Let X be a separable resolvable space. Then there exists at most countable set $A \subseteq X$ such that $\overline{A} = \overline{X \setminus A} = X$.*

Proof. Let X_1, X_2 and D be dense subsets of X such that $X = X_1 \sqcup X_2$ and $|D| \leq \aleph_0$. Consider the system

$$\mathcal{U} = \{U \in \mathcal{T}(X) : U \subseteq \overline{D \cap X_i} \text{ for an } i \in \{1,2\}\}.$$

We claim that $G = \cup\mathcal{U}$ is dense in X. Indeed, suppose that there exists a nonempty set $V \in \mathcal{T}(X)$ such that $V \cap G = \emptyset$. Then $X_i \cap D$ is nowhere dense in V for $i = 1, 2$, which implies that $D \cap V = (D \cap X_1 \cap V) \cup (D \cap X_2 \cap V)$ is nowhere dense in V. But $D \cap V$ is dense in V since D is dense in X, a contradiction.

Let $\mathcal{V}$ be a maximal subsystem of $\mathcal{U}$ which consists of disjoint elements. Note that $\cup\mathcal{V}$ is dense in G. For every $V \in \mathcal{V}$ there exists $i_V \in \{1,2\}$ such that $X_{i_V} \cap D$ is dense in V. Remark that $V \setminus (X_{i_V} \cap D)$ is dense in V too. Put $A = \bigsqcup_{V \in \mathcal{V}} (X_{i_V} \cap D \cap V)$. Since A and $X \setminus A$ are dense in G, we have that $\overline{A} = \overline{X \setminus A} = X$. Moreover, $|A| \leq \aleph_0$ since $A \subseteq D$. □

Proposition 3.13. *Let X be a separable resolvable Hausdorff space such that a completion to any countable set is a connected set, Y be a T_1-space, $|Y| \geq 2$. Then*

$$\mathrm{sG}(X,Y) \setminus \mathrm{PC}(X,Y) \neq \emptyset.$$

Proof. It follows from Lemma 3.12 that there exists at most countable set $A \subseteq X$ such that $\overline{A} = \overline{X \setminus A} = X$. Fix two distinct points $y_1, y_2 \in Y$ and consider a function $f : X \to Y$, $f(x) = y_1$ if $x \in A$, and $f(x) = y_2$ if $x \in X \setminus A$. According to Proposition 3.8, $f \in \mathrm{sG}(X,Y)$.

We will show that $f \notin \mathrm{PC}(X,Y)$. Fix $x \in A$ and let $V = Y \setminus \{y_2\}$. Since X is Hausdorff, there exists an open neighborhood U of x such that $X \setminus \overline{U} \neq \emptyset$. Assume that there exists an open neighborhood W of x such that $W \subseteq U$ and $f(\mathrm{fr}\, W) \subseteq V$. Then $\mathrm{fr}\, W \subseteq A$, which implies that $|\mathrm{fr}\, W| \leq \aleph_0$. Notice that $X \setminus \mathrm{fr}\, W = W \sqcup (X \setminus \overline{W})$. The last equality implies a contradiction, because $X \setminus \mathrm{fr}\, W$ is connected and $W \sqcup (X \setminus \overline{W})$ is not since W and $X \setminus \overline{W}$ are open nonempty sets. Thus, f is not peripherally continuous at x. □

4. On Gibson Baire-one functions

In this section we show that the Gibson property is equivalent to the almost continuity (in the sense of Husain) and prove that every Gibson Baire-one function from a Baire space to a metric space is continuous.

Definition 4.1. A function $f : X \to Y$ is called *almost continuous (in the sense of Husain)* if $f^{-1}(W) \subseteq \operatorname{int}\overline{f^{-1}(W)}$ for any $W \in \mathcal{T}(Y)$.

Proposition 4.2. *Let X and Y be topological spaces. Then $f : X \to Y$ is a Gibson function if and only if f is almost continuous.*

Proof. Necessity. Let $f : X \to Y$ be a Gibson function, $W \in \mathcal{T}(Y)$ and $x \in f^{-1}(W)$. Denote $G = X \setminus \overline{f^{-1}(W)}$ and assume that $x \in \overline{G}$. Since f is Gibson at x, $f(G) \cap W \neq \emptyset$. From the other side, it is easily seen that $f(G) \subseteq Y \setminus W$, which implies a contradiction.

Sufficiency. Choose any $G \in \mathcal{T}(X)$, $x \in \operatorname{fr} G$ and an open neighborhood W of $f(x)$ in Y. Denote $U = \operatorname{int}\overline{f^{-1}(W)}$. Since $x \in f^{-1}(W) \subseteq U$, $U \cap G \neq \emptyset$. Notice that $f^{-1}(W)$ is dense in $U \cap G$. Then there exists an $x' \in U \cap G$ such that $f(x') \in W$. Hence, f is Gibson at x. □

Proposition 4.3. *Let X be a topological space, Y be a regular space, $f : X \to Y$ be a Gibson function with $D(f) \neq \emptyset$. Then there exists an open nonempty set $U \subseteq X$ such that $D(f|_U) = U$.*

Proof. Suppose that f is discontinuous at a point x_0. Then there exists an open neighborhood W of $f(x_0)$ such that $V \cap f^{-1}(Y \setminus W) \neq \emptyset$ for any open neighborhood V of x_0 in X. It follows from the regularity of Y that there exists an open neighborhood W_1 of $f(x_0)$ such that $\overline{W}_1 \subseteq W$. Denote $W_2 = Y \setminus \overline{W}$, $G_i = \operatorname{int}\overline{f^{-1}(W_i)}$ for $i = 1, 2$ and prove that $U = G_1 \cap G_2$ is nonempty. Indeed, according to Proposition 4.2, the set G_1 is a neighborhood of x_0. Then $G_1 \cap f^{-1}(W_2) \neq \emptyset$. Moreover, Proposition 4.2 implies that $f^{-1}(W_2) \subseteq G_2$. Therefore, $U \supseteq G_1 \cap f^{-1}(W_2) \neq \emptyset$.

Now we show that $C(f|_U) = \emptyset$. Assume $f|_U$ is continuous at $x \in U$. If $f(x) \in W_1$ ($f(x) \in W_2$) then there exists a neighborhood V of x in U such that $f(V) \subseteq W_1$ ($f(V) \subseteq W_2$), which contradicts the density of $f^{-1}(W_2)$ ($f^{-1}(W_1)$) in U. If $f(x) \in \overline{W} \setminus W_1$ then there exist open neighborhoods W' of $f(x)$ and V of x in U such that $W' \cap W_1 = \emptyset$ and $f(V) \subseteq W'$. But $V \cap f^{-1}(W_1) \neq \emptyset$, a contradiction. Hence, $x \in D(f|_U)$. □

Taking into account that every Baire-one function $f : X \to Y$, where X is a Baire space and Y is a metric space, is pointwise discontinuous, the result below immediately follows from [1, Theorem 2.1] (see also [2, Corollary 1]). We present a proof here for convenience of the reader.

Theorem 4.4. *Let X be a Baire space and Y is a metric space. Then every Gibson Baire-one function $f : X \to Y$ is continuous.*

Proof. Assume there exists a Gibson function $f \in \mathrm{B}_1(X, Y)$ which is discontinuous. Then Proposition 4.3 implies that the restriction $f|_U$ on an open nonempty set $U \subseteq X$ is everywhere discontinuous. Furthermore, $f|_U$ is of the first Baire class and according to [13, p. 405] the set $D(f|_U)$ is of the first category in U. From the other hand, X is a Baire space. Then U is of the second category in itself, a contradiction. □

5. Segmentary connected functions

Let X be a vector space. For $a, b \in X$ we set

$$[a, b] = \{(1 - t)a + tb : t \in [0, 1]\}, \qquad [a, b) = \{(1 - t)a + tb : t \in [0, 1)\}.$$

The set $[a, b]$ is said to be *a segment in X*.

Definition 5.1. *Let Y be a topological space. A function $f : X \to Y$ is called* `segmentary connected`*, $f \in \mathrm{S}(X, Y)$, if $f([a, b])$ is a connected set for any segment $[a, b] \subseteq X$.*

The definition immediately implies that $\mathrm{D}(X, Y) \subseteq \mathrm{S}(X, Y)$ for a topological vector space X and for a topological space Y. It is not difficult to verify that $\mathrm{D}(X, Y) = \mathrm{S}(X, Y)$ if X is a connected subset of $\mathbb{R}$ and Y is a topological space.

Proposition 5.2. $\mathrm{S}(\mathbb{R}^2, [0, 1]) \setminus \mathrm{wG}(\mathbb{R}^2, [0, 1]) \neq \emptyset$.

Proof. Consider two logarithmic spirals $S_1 \subseteq \mathbb{R}^2$ and $S_2 \subseteq \mathbb{R}^2$, which in polar coordinates can be written as $\varrho = e^{\varphi}$ and $\varrho = 2e^{\varphi}$, respectively. Set $p_0 = (0, 0)$. Since S_1 and S_2 are closed in $X = \mathbb{R}^2 \setminus \{p_0\}$ and disjoint, there exists such a continuous function $\alpha : X \to [0, 1]$ that $S_i = \alpha^{-1}(i-1)$, $i = 1, 2$. Let $f(p) = \alpha(p)$ if $p \neq p_0$, and $f(p_0) = 1$.

We prove that $f \in \mathrm{S}(\mathbb{R}^2, [0, 1])$. Indeed, if $[a, b]$ is such a segment in $\mathbb{R}^2$ that $p_0 \notin [a, b]$, then the restriction $f|_{[a,b]}$ is continuous, consequently, $f([a, b])$ is a connected set. Now let $p_0 \in [a, b]$. It is obvious that there exist $p_1 \in S_1 \cap [a, b]$ and $p_2 \in S_2 \cap [a, b]$. Since $f(p_1) = 0$, $f(p_2) = 1$ and $f|_{[p_1,p_2]}$ is continuous, $f([p_1, p_2]) = [0, 1]$. Taking unto account that $f([p_1, p_2]) \subseteq f([a, b]) \subseteq [0, 1]$, we have $f([a, b]) = [0, 1]$. Thus, f is segmentary connected.

We show now that $f \notin \mathrm{wG}(\mathbb{R}^2, [0, 1])$. Let $G_0 = f^{-1}([0, \frac{1}{2}))$. Choose any point $p \in S_1$ and consider a component G of p in G_0. Then $G \in \mathcal{G}(\mathbb{R}^2)$. Since $p_0 \in \overline{S}_1$ and $S_1 \subseteq G$, $p_0 \in \overline{G}$. But $f(p_0) = 1 \notin \overline{f(G)} \subseteq [0, \frac{1}{2}]$. Therefore, f does not satisfy the weak Gibson property at p_0. □

We will need the following two auxiliary facts.

Lemma 5.3. *Let X be a topological space, $f : X \to Y$ be a function, $A \subseteq X$, $A = \bigcup_{i\in I} A_i$, where $f(A_i)$ is connected for every $i \in I$, and for each two points $x_1, x_2 \in A$ there exists a finite set of indexes $i_1, \dots, i_n \in I$ such that $x_1 \in A_{i_1}$, $x_2 \in A_{i_n}$ and $A_{i_k} \cap A_{i_{k+1}} \neq \emptyset$ for $1 \leq k < n$. Then $f(A)$ is a connected set.*

Proof. It is easy to see that each two points $y_1, y_2 \in f(A)$ are contained in a connected subset of $f(A)$. Thus, $f(A)$ is a connected set by [14, p. 141]. □

Lemma 5.4. *Let X be a locally connected space, $\mathcal{B}$ be a base of open connected sets in X, Y be a topological space and $f : X \to Y$. If $f(U)$ is connected for each $U \in \mathcal{B}$, then $f \in \mathrm{wD}(X, Y)$.*

Proof. Choose an arbitrary set $G \in \mathcal{G}(X)$. Then $G = \bigcup_{i\in I} B_i$, where $B_i \in \mathcal{B}$. According to [14, p. 144] the sets B_i for $i \in I$ satisfy the condition of Lemma 5.3. Thus, $f(G)$ is connected. □

Proposition 5.5. *Let X be a topological vector space, Y be a topological space. Then $\mathrm{S}(X, Y) \subseteq \mathrm{wD}(X, Y)$.*

Proof. Let $f \in \mathrm{S}(X, Y)$. Choose a base $\mathcal{U}$ of balanced open neighborhoods of zero and write $\mathcal{B} = (U + x : x \in X, U \in \mathcal{U})$. Then $\mathcal{B}$ is a base of open connected sets in X. Take an arbitrary $B \in \mathcal{B}$. Then $B = x + U$ for some $x \in X$ and $U \in \mathcal{U}$. Since U is balanced, $[x, y] \subseteq B$ for each $y \in B$, consequently, $B = \bigcup_{y\in B} [x, y]$. Then $f(B) = \bigcup_{y\in B} f([x, y])$. Since $f([x, y])$ is connected for all $y \in B$ and $\bigcap_{y\in B} f([x, y]) \neq \emptyset$, $f(B)$ is connected too. Hence, $f \in \mathrm{wD}(X, Y)$ by Proposition 5.4. □

Remark that Proposition 3.9 implies that $\mathrm{wG}(\mathbb{R}, Y) \setminus \mathrm{S}(\mathbb{R}, Y) \neq \emptyset$ for every T_1-space Y with $|Y| \geq 2$. However, the following result shows that if a function $f \in \mathrm{wG}(X, Y)$ has "many" points of continuity, then it is segmentary connected.

Theorem 5.6. *Let X be a topological vector Hausdorff space, Y be a topological space and let $f \in \mathrm{wG}(X, Y)$ be such a function that $D(f) \cap \ell$ is locally finite for every line $\ell \in X$. Then $f \in \mathrm{S}(X, Y)$.*

Proof. Fix an arbitrary segment $S = [a, b] \subseteq X$.

We first consider the case $|D(f) \cap S| \leq 1$. If $f|_S$ is continuous, then, evidently, $f(S)$ is connected. Now let $D(f) \cap S = \{x_0\}$. Suppose that $f(S)$ is not connected. Then at least one of the sets $f([a, x_0])$ or $f([x_0, b])$ is not connected. Without loss the generality we may assume that $f([a, x_0])$ is not connected. Since $f|_{[a,x_0)}$ is continuous, $f([a, x_0))$ is connected. Then $f(x_0) \notin \overline{f([a, x_0))}$. Since X is a topological vector Hausdorff space, it is regular, consequently, there are open disjoint sets W_1 and W_2 in Y such that $f(x_0) \in W_1$ and $f([a, x_0)) \in W_2$. The

continuity of f on $S\setminus\{x_0\}$ implies that for every $x\in[a,x_0)$ there exists an open connected neighborhood U_x of x in S such that $x_0\notin U_x$ and $f(U_x)\subseteq W_2$. It is easy to check that $V=\bigcup\limits_{x\in[a,x_0)}U_x$ belong to $\mathcal{G}(X)$ and $x_0\in\overline{V}$. Then, since $f\in \mathrm{wG}(X,Y)$, we have $f(x_0)\subseteq\overline{f(V)}\subseteq\overline{W_2}$, which implies a contradiction, because $W_1\cap\overline{W_2}=\emptyset$.

Now consider the general case. Then for every $x\in S$ there exist $a_x,b_x\in S$ such that $|D(f)\cap[a_x,b_x]|\leq 1$. Let $S_x=[a_x,b_x]$. Then $S=\bigcup\limits_{x\in S}S_x$. Moreover, it follows from what has already been proved that $f(S_x)$ is connected for every $x\in S$. Combining [14, p. 144] and Lemma 5.3 gives the connectedness of $f(S)$. □

6. Baire-one functions with connected graphs

Theorem 6.1. *Let X be a connected and locally connected space, Y be a topological space and let $f:X\to Y$ be a weakly Gibson barely continuous function. Then*

(1) $f\in \mathrm{CG}(X,Y)$;
(2) $f\in \mathrm{wD}(X,Y)$.

Proof. **(1).** Suppose, contrary to our claim, that $\mathrm{Gr}(f)=\Gamma_1\sqcup\Gamma_2$, where Γ_1 and Γ_2 are closed and open in $\mathrm{Gr}(f)$. For $i=1,2$ write $X_i=\{x\in X:(x,f(x))\in\Gamma_i\}$. Clearly, $X=X_1\sqcup X_2$. As X is connected we have $F=\overline{X}_1\cap\overline{X}_2$ is nonempty. Since f is barely continuous, there exists $x_0\in C(f|_F)$. We may assume that $x_0\in X_1$. Since Γ_1 is open in $\mathrm{Gr}(f)$, there exist open neighborhoods U and V of x_0 and $f(x_0)$ in X and Y, respectively, such that $(U\times V)\cap\mathrm{Gr}(f)\subseteq\Gamma_1$. Notice that the continuity of $f|_F$ at x_0 and the locally connectedness of X imply that we can choose U such that $f(U\cap F)\subseteq V$ and U is connected. Then $U\cap F\subseteq X_1$ since $\mathrm{Gr}(f|_{U\cap F})\subseteq U\times V$.

Since $x_0\in\overline{X}_2$, $U\cap X_2\neq\emptyset$. Choose any $x'\in U\cap X_2$. Then $x'\notin F$. Let G be a component of x' in $U\setminus\overline{X}_1$. Notice that $U\cap G\neq\emptyset\neq U\setminus G$. Then, $U\cap\mathrm{fr}G\neq\emptyset$ by Lemma 3.10. It is not difficult to verify that $U\cap\mathrm{fr}G\subseteq F$ since G is a component. Fix $x''\in U\cap\mathrm{fr}G$. Then $f(x'')\in V$. But $f\in\mathrm{wG}(X,Y)$ and $x''\in\overline{G}$, then $f(x'')\in\overline{f(G)}\subseteq\overline{Y\setminus V}=Y\setminus V$ since $f(G)\subseteq Y\setminus V$, a contradiction.

(2). Let $G\in\mathcal{G}(X)$. Obviously, the restriction $f|_G$ has the weak Gibson property and is barely continuous on G. Since G is open in X, it is locally connected. Then (1) implies that $\mathrm{Gr}(f|_G)$ is connected in $G\times f(G)$. If $f(G)=B_1\sqcup B_2$, where B_i is open and closed in $f(G)$ for $i=1,2$, then $\mathrm{Gr}(f|_G)\subseteq(G\times B_1)\sqcup(G\times B_2)$, where $G\times B_i$ is open and closed in $G\times f(G)$ for $i=1,2$, which contradicts the connectedness of $\mathrm{Gr}(f|_G)$. □

Theorem 6.2. *Let X be a topological space, Y be a T_1-space and $f : X \to Y$ be a barely continuous function. If*

(1) *X is arcwise connected and f is a Darboux function, or*
(2) *X is a topological vector space and f is segmentary connected,*

then $f \in \mathrm{CG}(X, Y)$.

Proof. Fix $x_0 \in X$. In the case (1) for every point $x \in X$, $x \neq x_0$, there exists a homeomorphic embedding $h_x : [0, 1] \to X$ satisfying $h_x(0) = x_0$ and $h_x(1) = x$. Let $X_0 = X \setminus \{x_0\}$ and $A_x = h_x([0, 1])$ in the case (1) (and $A_x = [x_0, x]$ in the case (2)) for all $x \in X_0$. Then $X = \bigcup\limits_{x \in X_0} A_x$. For every $x \in X_0$ we consider a function $g_x = f|_{A_x}$. Then $g_x : A_x \to Y$ is pointwise discontinuous and $g_x \in \mathrm{D}(A_x, Y)$. By Proposition 3.4 $g_x \in \mathrm{wG}(A_x, Y)$.

Since A_x is homeomorphic to $[0, 1]$, it is connected and locally connected. Then Theorem 6.1(1) implies that $g_x \in \mathrm{CG}(A_x, Y)$ for all $x \in X_0$. Notice that $\mathrm{Gr}(f) = \bigcup\limits_{x \in X_0} \mathrm{Gr}(g_x)$. For every $x \in X_0$ the set $\mathrm{Gr}(g_x)$ is connected. Moreover, $(x_0, f(x_0)) \in \bigcap\limits_{x \in X_0} \mathrm{Gr}(g_x)$. Hence, $\mathrm{Gr}(f)$ is a connected set. □

Since every Baire-one function between a hereditarily Baire space and a metric space is barely continuous [3, p. 125], we have the following corollary from Theorem 6.1 and Theorem 6.2.

Theorem 6.3. *Let X be a hereditarily Baire space, Y be a metric space and $f \in \mathrm{B}_1(X, Y)$. If*

(1) *X is a connected and locally connected space and $f \in \mathrm{wG}(X, Y)$, or*
(2) *X is an arcwise connected space and $f \in \mathrm{D}(X, Y)$, or*
(3) *X is a topological vector space and $f \in \mathrm{S}(X, Y)$,*

then f has a connected graph.

Remark that a function $f \in \mathrm{B}_1(\mathbb{R}^2, \mathbb{R})$ with a connected graph does not necessarily have either the weak Darboux property, or the weak Gibson property. Indeed, if we consider a function $f : \mathbb{R}^2 \to \mathbb{R}$, $f(x, 0) = -x^2$ for $|x| \leq 1$ and $f(x, y) = 0$ for $(x, y) \in \mathbb{R}^2 \setminus ([-1, 1] \times \{0\})$, then f satisfy the required conditions.

If $X = Y = \mathbb{R}$, then the conditions (1), (2) and (3) of Theorem 6.3 are equivalent. If $X = \mathbb{R}^n$ $(n \geq 2)$, $Y = \mathbb{R}$, then (2) $\Rightarrow$ (1) and (2) $\Rightarrow$ (3). Example 5.2 gives (3) $\not\Rightarrow$ (1) and, consequently, (3) $\not\Rightarrow$ (2). Proposition 6.4 below shows, in particular, that (1) $\not\Rightarrow$ (3) and, consequently, (1) $\not\Rightarrow$ (2).

Proposition 6.4. $(\mathrm{sG}(\mathbb{R}^2, \mathbb{R}) \cap B_1(\mathbb{R}^2, \mathbb{R})) \setminus \mathrm{S}(\mathbb{R}^2, \mathbb{R}) \neq \emptyset$.

Proof. Let $(P_n)_{n=1}^{\infty}$ be a sequence of rectangles $P_n = (a_n, b_n) \times (c_n, d_n)$ with the following properties:

(1) $\overline{P}_n \cap \{(x, 0) : x \in \mathbb{R}\} = \emptyset$ for every $n \in \mathbb{N}$;

(2) $\overline{P}_n \cap \overline{P}_m = \emptyset$ for distinct $m, n \in \mathbb{N}$;
(3) $A = \bigcup\limits_{n=1}^{\infty} P_n$ is dense in $\mathbb{R}^2$;
(4) $\lim\limits_{n\to\infty} \operatorname{diam} P_n = 0$.

We will show that for every nonempty set $G \in \mathcal{G}(\mathbb{R}^2)$ there exists such an $n \in \mathbb{N}$ that either $P_n \subseteq G$, or $G \subseteq P_n$. Indeed, fix any nonempty open connected set $G \subseteq \mathbb{R}^2$. If $G \subseteq A$ then choosing $n \in \mathbb{N}$ such that $G \cap P_n \neq \emptyset$, we have $G \subseteq P_n$ since G is connected. Now denote $B = \mathbb{R}^2 \setminus A$ and assume that $G \cap B \neq \emptyset$. Take $p_0 \in G \cap B$ and $\delta > 0$ such that $K_1 = \{p \in \mathbb{R}^2 : d(p, p_0) < 2\delta\} \subseteq G$. Let $N_1 = \{n \in \mathbb{N} : p_0 \notin \overline{P}_n \text{ and } \operatorname{diam} P_n > \delta\}$. Then $|N_1| < \aleph_0$ and there exists $\delta_1 < \delta$ such that $K_2 \cap (\bigcup\limits_{n\in N_1} \overline{P}_n) = \emptyset$, where $K_2 = \{p \in \mathbb{R}^2 : d(p, p_0) < \delta_1\}$. Denote $N_2 = \{n \in \mathbb{N} : p_0 \in \overline{P}_n\}$. Remark that $|N_2| \leq 1$ according to (2). Then, taking into account that $p_0 \notin \bigcup\limits_{n\in N_2} P_n$, we have $G_1 = K_2 \setminus (\bigcup\limits_{n\in N_2} \overline{P}_n)$ is nonempty set. Since B is nowhere dense in $\mathbb{R}^2$, there exists such a number n that $P_n \cap G_1 \neq \emptyset$. Since $n \notin N_1 \cup N_2$, $\operatorname{diam} P_n \leq \delta$. Choose $p_1 \in P_n \cap G_1$. Then for every $p \in P_n$

$$d(p, p_0) \leq d(p, p_1) + d(p_1, p_0) \leq \delta + \delta_1 < 2\delta.$$

Therefore, $P_n \subseteq K_1 \subseteq G$.

For every $n \in \mathbb{N}$ we consider a continuous function $f_n : \overline{P}_n \to [0, 1]$ such that $f_n(P_n) = (0, 1]$ and $f_n(\operatorname{fr} P_n) = \{0\}$. Define a function $f : \mathbb{R}^2 \to \mathbb{R}$ by $f(p) = f_n(p)$ if $p \in P_n$, $f(0, 0) = 1$ and $f(p) = 0$ if $p \in B \setminus \{0, 0\}$. Notice that the restriction $f|_A$ is continuous and the restriction $f|_B$ is of the first Baire class. Hence, $f \in B_1(\mathbb{R}^2, \mathbb{R})$. Moreover, for the segment $S = \{(x, 0) : x \in [0, 1]\}$ we have $f(S) = \{0, 1\}$. Thus, $f \notin \mathrm{S}(\mathbb{R}^2, \mathbb{R})$.

Finally, we will prove that $f \in \mathrm{sG}(\mathbb{R}^2, \mathbb{R})$. Let $p_0 \in \mathbb{R}^2$, $G \in \mathcal{G}(\mathbb{R}^2)$, $p_0 \in \overline{G}$ and $U \subseteq \mathbb{R}^2$ is an open neighborhood of p_0. If $p_0 \in A$ then the continuity of f at p_0 implies that $f(p_0) \in \overline{f(U \cap G)}$. Now assume $p_0 \in B$. If there exists such an $n \in \mathbb{N}$ that $p_0 \in \overline{G \cap P_n}$, then $p_0 \in \overline{U \cap G \cap P_n}$ and the continuity of $f|_{\overline{P}_n}$ implies that $f(p_0) \in \overline{f(U \cap G \cap P_n)}$, besides, $f(p_0) \in \overline{f(U \cap G)}$.

We will show that there is $p_1 \in U \cap G \cap B$ if $p_0 \notin \overline{G \cap P_n}$ for every $n \in \mathbb{N}$. Choose $\delta > 0$ such that $\{p \in \mathbb{R}^2 : d(p, p_0) < 2\delta\} \subseteq U$. Since $p_0 \notin \overline{G \cap P_n}$ for all $n \in \mathbb{N}$, it follows from (4) that $p_0 \notin \overline{\bigcup\limits_{\operatorname{diam}(P_n)>\delta} G \cap P_n}$. Taking into account that $p_0 \in \overline{G}$, we obtain that the set $\{k \in \mathbb{N} : G \cap P_n \cap U \neq \emptyset\}$ is infinite. Take $n \in \mathbb{N}$ such that $\operatorname{diam} P_n < \delta$ and there is $p \in P_n \cap G$ with $d(p, p_0) < \delta$. Then $\overline{P}_n \subseteq U$. Notice that $P_n \cap G$ is open nonempty subset of the connected set G which is not coincide with G. Then $P_n \cap G$ is not closed in G. Therefore, there exists $p_1 \in \operatorname{fr} P_n \cap G \subseteq U \cap G \cap B$.

Let G_1 is a component of p_1 in $G \cap U$. Since $p_1 \notin A$, $G_1 \not\subseteq P_m$ for every $m \in \mathbb{N}$. Thus, from what has already been proved, it follows that there exists such a number m that $P_m \subseteq G_1$. Then

$$f(p_0) \in [0,1] = \overline{f(P_m)} \subseteq \overline{f(G_1)} \subseteq \overline{f(G \cap U)}.$$

Hence, f is strongly Gibson. □

References

[1] BERNER, A.J.: *Almost continuous functions with closed graphs*, Canad. Math. Bull. **25** (1982), 428–434.

[2] BORSÍK, J.—DOBOŠ, J.: *On certain decompositions of continuity*, Rend. Ist. Mat. Univ. Trieste **20** (1988), 275–282.

[3] BOURBAKI, N.: *General Topology*, M.: Nauka, 1975 (in Russian).

[4] CIESIELSKI, K.—JASTRZĘBSKI, J.: *Darboux-like functions whithin the classes of Baire-one, Baire-two and additive functions* // Topology Appl., **103**(2000), 203–219.

[5] Encyclopedia of General Topology [ed. by K.Hart, J.Nagata, J.Vaughan], 2004.

[6] ENGELKING, R.: *General Topology*, Revised and completed edition, Heldermann Verlag, Berlin, 1989.

[7] EVANS, M.J.—HUMKE, P.D.: *Baire one, Gibson and weakly Gibson real functions of several variables*, Rend. Circ. Mat. Palermo **59** (2010), 47–51.

[8] FARKOVÁ, J.: *Note on a theorem of Kuratowski-Sierpiński*, Matematický časopis, **23**(2) (1973), 179–183.

[9] GIBSON, R.—NATKANIEC, T.: *Darboux-like functions*, Real Anal. Exchange **22**(2) (1996-97), 492–533.

[10] HAGAN, M. R.: *Equivalence of connectivity maps and peripherally continuous transformations*, Proc. Amer. Math. Soc. **17** (1966), 175–177.

[11] KELLUM, K.: *Functions that separate* $X \times \mathbb{R}$, Houston J. Math. **36** (2010), 1221–1226.

[12] KURATOWSKI, C.—SIERPINSKI, W.: *Les fonctions de classe I et les ensembles connexes punctiformes*, Fund. Math. 3 (1922), 303–313.

[13] KURATOWSKI, K.: *Topology*, Volume 1, Moscow, 1966 (in Russian).

[14] KURATOWSKI, K.: *Topology*, Volume 2, Moscow, 1969 (in Russian).

[15] WHYBURN, G. T.: *Connectivity of peripherally continuous functions*, Proc. Nat. Acad. Sci. USA **55** (1966), 1040–1041.

[16] YOUNG, J.: *A theorem in the theory of functions of a real variable*, Rend. Circ. Mat. Palermo **24** (1907), 187–192.

* Chernivtsi National University
Department of Mathematical Analysis
Kotsjubyns'koho 2
Chernivtsi 58012
UKRAINE
E-mail address: Maslenizza.Ua@gmail.com

** Chernivtsi National University
Department of Mathematical Analysis
Kotsjubyns'koho 2
Chernivtsi 58012
UKRAINE
E-mail address: vmykhaylyuk@ukr.net